

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

## **1. NAME OF THE MEDICINAL PRODUCT**

Polivy 30 mg powder for concentrate for solution for infusion.  
Polivy 140 mg powder for concentrate for solution for infusion.

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

### Polivy 30 mg powder for concentrate for solution for infusion

Each vial of powder for concentrate for solution for infusion contains 30 mg of polatuzumab vedotin.

After reconstitution, each mL contains 20 mg of polatuzumab vedotin.

### Polivy 140 mg powder for concentrate for solution for infusion

Each vial of powder for concentrate for solution for infusion contains 140 mg of polatuzumab vedotin.

After reconstitution, each mL contains 20 mg of polatuzumab vedotin.

Polatuzumab vedotin is an antibody-drug conjugate composed of the anti-mitotic agent monomethyl auristatin E (MMAE) covalently conjugated to a CD79b-directed monoclonal antibody (recombinant humanized immunoglobulin G1 [IgG1], produced in Chinese Hamster Ovary cells by recombinant DNA technology).

For the full list of excipients, see section 6.1.

## **3. PHARMACEUTICAL FORM**

Powder for concentrate for solution for infusion (powder for concentrate).

White to greyish-white lyophilized cake.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Polivy in combination with rituximab, cyclophosphamide, doxorubicin, and prednisone (R-CHP) is indicated for the treatment of adult patients with previously untreated diffuse large B-cell lymphoma (DLBCL).

Polivy in combination with bendamustine and rituximab is indicated for the treatment of adult patients with relapsed/refractory diffuse large B-cell lymphoma (DLBCL) who are not candidates for haematopoietic stem cell transplant.

### **4.2 Posology and method of administration**

Polivy must only be administered under the supervision of a healthcare professional experienced in the diagnosis and treatment of cancer patients.

## Posology

### **Diffuse large B-cell lymphoma**

#### *Previously untreated patients*

The recommended dose of Polivy is 1.8 mg/kg, given as an intravenous infusion every 21 days in combination with rituximab, cyclophosphamide, doxorubicin, and prednisone (R-CHP) for 6 cycles. Polivy, rituximab, cyclophosphamide and doxorubicin can be administered in any order on Day 1 after the administration of prednisone. Prednisone is administered on Days 1-5 of each cycle. Cycles 7 and 8 consist of rituximab as monotherapy.

Refer to the summary of product characteristics (SmPC) of chemotherapy agents given in combination with Polivy for patients with previously untreated DLBCL.

#### *Relapsed or refractory patients*

The recommended dose of Polivy is 1.8 mg/kg, given as an intravenous infusion every 21 days in combination with bendamustine and rituximab for 6 cycles. Polivy, bendamustine and rituximab can be administered in any order on Day 1 of each cycle. When administered with Polivy, the recommended dose of bendamustine is 90 mg/m<sup>2</sup>/day on Day 1 and Day 2 of each cycle and the recommended dose of rituximab is 375 mg/m<sup>2</sup> on Day 1 of each cycle. Due to limited clinical experience in patients treated with 1.8 mg/kg Polivy at a total dose >240 mg, it is recommended not to exceed the dose 240 mg/cycle.

#### *Previously untreated and relapsed or refractory patients*

If not already premedicated, premedication with an antihistamine and anti-pyretic should be administered to patients prior to Polivy.

#### *Delayed or missed doses*

If a planned dose of Polivy is missed, it should be administered as soon as possible and the schedule of administration should be adjusted to maintain a 21-day interval between doses.

#### *Dose modifications*

The infusion rate of Polivy should be slowed or interrupted if the patient develops an infusion-related reaction. Polivy should be discontinued immediately and permanently if the patient experiences a life-threatening reaction.

There are different potential dose modifications for Polivy in patients with previously untreated DLBCL and those who are relapsed or refractory.

For dose modifications to manage peripheral neuropathy (section 4.4) see Table 1 below.

**Table 1 Polivy dose modifications for peripheral neuropathy (PN)**

| <b>Indication</b>          | <b>Severity of PN on Day 1 of any cycle</b> | <b>Dose modification</b>   |
|----------------------------|---|--|
| Previously untreated DLBCL | Grade 2 <sup>a</sup>                        | <p>Sensory neuropathy:</p> <ul style="list-style-type: none"> <li>• Reduce Polivy to 1.4 mg/kg.</li> <li>• If Grade 2 persists or recurs at Day 1 of a future cycle, reduce Polivy to 1.0 mg/kg.</li> <li>• If already at 1.0 mg/kg and Grade 2 occurs at Day 1 of a future cycle, discontinue Polivy.</li> </ul> <p>Motor neuropathy:</p> <ul style="list-style-type: none"> <li>• Withhold Polivy dosing until improvement to Grade <math>\leq</math>1.</li> <li>• Restart Polivy at the next cycle at 1.4 mg/kg.</li> <li>• If already at 1.4 mg/kg and Grade 2 occurs at Day 1 of a future cycle, withhold Polivy dosing until improvement to Grade <math>\leq</math> 1. Restart Polivy at 1.0 mg/kg.</li> <li>• If already at 1.0 mg/kg and Grade 2 occurs at Day 1 of a future cycle, discontinue Polivy.</li> </ul> <p>If concurrent sensory and motor neuropathy, follow the most severe restriction recommendation above.</p> |
|                            | Grade 3 <sup>a</sup>                        | <p>Sensory neuropathy:</p> <ul style="list-style-type: none"> <li>• Withhold Polivy dosing until improvement to Grade <math>\leq</math> 2.</li> <li>• Reduce Polivy to 1.4 mg/kg.</li> <li>• If already at 1.4 mg/kg, reduce Polivy to 1.0 mg/kg. If already at 1.0 mg/kg, discontinue Polivy.</li> </ul> <p>Motor neuropathy:</p> <ul style="list-style-type: none"> <li>• Withhold Polivy dosing until improvement to Grade <math>\leq</math> 1.</li> <li>• Restart Polivy at the next cycle at 1.4 mg/kg.</li> <li>• If already at 1.4 mg/kg and Grade 2–3 occurs, withhold Polivy dosing until improvement to Grade <math>\leq</math> 1. Restart Polivy at 1.0 mg/kg.</li> <li>• If already at 1.0 mg/kg and Grade 2–3 occurs, discontinue Polivy.</li> </ul> <p>If concurrent sensory and motor neuropathy, follow the most severe restriction recommendation above.</p>  |
|                            | Grade 4                                     | Discontinue Polivy.  |
| R/R DLBCL                  | Grade 2–3                                   | <p>Withhold Polivy dosing until improvement to <math>\leq</math> Grade 1.</p> <p>If recovered to Grade <math>\leq</math> 1 on or before Day 14, restart Polivy at a permanently reduced dose of 1.4 mg/kg.</p> <p>If a prior dose reduction to 1.4 mg/kg has occurred, discontinue Polivy.</p> <p>If not recovered to Grade <math>\leq</math> 1 on or before Day 14, discontinue Polivy.</p>   |
|                            | Grade 4                                     | Discontinue Polivy.  |

<sup>a</sup> R-CHP may continue to be administered.

For dose modifications to manage myelosuppression (section 4.4) see Table 2 below.

**Table 2 Polivy, chemotherapy and rituximab dose modifications to manage myelosuppression**

| <b>Indication</b>          | <b>Severity of myelosuppression on Day 1 of any cycle</b> | <b>Dose modification</b>   |
|----------------------------|---|--|
| Previously untreated DLBCL | Grade 3–4 Neutropenia                                     | Withhold all treatment until ANC* recovers to > 1000/μL.<br>If ANC recovers to > 1000/μL on or before Day 7, resume all treatment without any dose reductions.<br>If ANC recovers to > 1000/μL after Day 7: <ul style="list-style-type: none"> <li>• resume all treatment; consider a dose reduction of cyclophosphamide and/or doxorubicin by 25-50%.</li> <li>• if cyclophosphamide and/or doxorubicin are already reduced by 25%, consider reducing one or both agents to 50%.</li> </ul>   |
|                            | Grade 3–4 Thrombocytopenia                                | Withhold all treatment until platelets recover to > 75,000/μL.<br>If platelets recover to > 75,000/μL on or before Day 7, resume all treatment without any dose reductions.<br>If platelets recover to > 75,000/μL after Day 7: <ul style="list-style-type: none"> <li>• resume all treatment; consider a dose reduction of cyclophosphamide and/or doxorubicin by 25-50%.</li> <li>• if cyclophosphamide and/or doxorubicin are already reduced by 25%, consider reducing one or both agents to 50%.</li> </ul>   |
| R/R DLBCL                  | Grade 3–4 Neutropenia <sup>1</sup>                        | Withhold all treatment until ANC recovers to > 1000/μL.<br>If ANC recovers to > 1000/μL on or before Day 7, resume all treatment without any additional dose reductions.<br>If ANC recovers to > 1000/μL after Day 7: <ul style="list-style-type: none"> <li>• restart all treatment with a dose reduction of bendamustine from 90 mg/m<sup>2</sup> to 70 mg/m<sup>2</sup> or 70 mg/m<sup>2</sup> to 50 mg/m<sup>2</sup>.</li> <li>• if a bendamustine dose reduction to 50 mg/m<sup>2</sup> has already occurred, discontinue all treatment.</li> </ul>           |
|                            | Grade 3–4 Thrombocytopenia <sup>1</sup>                   | Withhold all treatment until platelets recover to > 75,000/μL.<br>If platelets recover to > 75,000/μL on or before Day 7, resume all treatment without any dose reductions.<br>If platelets recover to > 75,000/μL after Day 7: <ul style="list-style-type: none"> <li>• restart all treatment with a dose reduction of bendamustine from 90 mg/m<sup>2</sup> to 70 mg/m<sup>2</sup> or 70 mg/m<sup>2</sup> to 50 mg/m<sup>2</sup>.</li> <li>• if a bendamustine dose reduction to 50 mg/m<sup>2</sup> has already occurred, discontinue all treatment.</li> </ul> |

<sup>1</sup>If primary cause is due to lymphoma, the dose of bendamustine may not need to be reduced.

\*ANC: absolute neutrophil count

For dose modifications to manage Infusion-related reactions (section 4.4) see Table 3 below.

**Table 3 Polivy dose modifications for Infusion-related reactions (IRRs)**

| Indication                         | Severity of IRR on Day 1 of any cycle | Dose modification  |
|------------------------------------|---------------------------------------|--|
| Previously untreated and R/R DLBCL | Grade 1–3 IRR                         | <p>Interrupt Polivy infusion and give supportive treatment.</p> <p>For the first instance of Grade 3 wheezing, bronchospasm, or generalized urticaria, permanently discontinue Polivy.</p> <p>For recurrent Grade 2 wheezing or urticaria, or for recurrence of any Grade 3 symptoms, permanently discontinue Polivy.</p> <p>Otherwise, upon complete resolution of symptoms, infusion may be resumed at 50% of the rate achieved prior to interruption. In the absence of infusion-related symptoms, the rate of infusion may be escalated in increments of 50 mg/hour every 30 minutes.</p> <p>For the next cycle, infuse Polivy over 90 minutes. If no infusion-related reaction occurs, subsequent infusions may be administered over 30 minutes. Administer premedication for all cycles.</p> |
|                                    | Grade 4 IRR                           | <p>Stop Polivy infusion immediately.</p> <p>Give supportive treatment.</p> <p>Permanently discontinue Polivy.</p>  |

### Special populations

#### *Elderly*

No dose adjustment of Polivy is required in patients  $\geq 65$  years of age (see section 5.2).

#### *Renal impairment*

No dose adjustment of Polivy is required in patients with creatinine clearance (CrCL)  $\geq 30$  mL/min. A recommended dose has not been determined for patients with CrCL  $< 30$  mL/min due to limited data.

#### *Hepatic impairment*

The administration of Polivy in patients with moderate or severe hepatic impairment (bilirubin greater than  $1.5 \times$  upper limit of normal [ULN]) should be avoided.

No adjustment in the starting dose is required when administering Polivy to patients with mild hepatic impairment (bilirubin greater than ULN to less than or equal to  $1.5 \times$  ULN or aspartate transaminase [AST] greater than ULN).

Per studied population in mild hepatic impairment (defined as AST or ALT  $> 1.0$  to  $2.5 \times$  ULN or total bilirubin  $> 1.0$  to  $1.5 \times$  ULN), there was a not more than 40% increase in unconjugated MMAE exposure, which was not deemed clinically significant.

#### *Paediatric population*

The safety and efficacy in children and adolescents less than 18 years have not been established. No data are available.

### Method of administration

Polivy is for intravenous use.

The initial dose of Polivy should be administered as a 90-minute intravenous infusion. Patients should be monitored for IRRs/hypersensitivity reactions during the infusion and for at least 90 minutes following completion of the initial dose.

If the prior infusion was well tolerated, the subsequent dose of Polivy may be administered as a 30-minute infusion and patients should be monitored during the infusion and for at least 30 minutes after completion of the infusion.

Polivy must be reconstituted and diluted using aseptic technique under the supervision of a healthcare professional. It should be administered as an intravenous infusion through a dedicated infusion line equipped with a sterile, non-pyrogenic, low-protein binding in-line or add-on filter (0.2 or 0.22 micrometer pore size) and catheter. Polivy must not be administered as intravenous push or bolus.

For instructions on reconstitution and dilution of the medicinal product before administration, see section 6.6.

### *Precaution to be taken before handling or administering the product*

Polivy contains a cytotoxic component which is covalently attached to the monoclonal antibody. Follow applicable proper handling and disposal procedure (see section 6.6).

## **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.  
Active severe infections (see section 4.4).

## **4.4 Special warnings and precautions for use**

### Traceability

In order to improve traceability of biological medicinal products, the trade name and the batch number of the administered product should be clearly recorded.

### Myelosuppression

Serious and severe neutropenia and febrile neutropenia have been reported in patients treated with Polivy as early as the first cycle of treatment. Prophylactic granulocyte colony stimulating factor (G-CSF) administration was required in the clinical development and should be considered. Grade 3 or 4 thrombocytopenia or anaemia can also occur with Polivy. Complete blood counts should be monitored prior to each dose of Polivy. More frequent lab monitoring and/or Polivy delays or discontinuation should be considered for patients with Grade 3 or Grade 4 neutropenia and/or thrombocytopenia (see section 4.2).

### Peripheral neuropathy (PN)

PN has been reported in patients treated with Polivy as early as the first cycle of treatment, and the risk increases with sequential doses. Patients with pre-existing PN may experience worsening of this condition. PN reported with treatment with Polivy is predominantly sensory PN. However, motor and sensorimotor PN have also been reported. Patients should be monitored for symptoms of PN such as hypoesthesia, hyperesthesia, paraesthesia, dysesthesia, neuropathic pain, burning sensation, muscle weakness, or gait disturbance. Patients experiencing new or worsening PN may require a delay, dose reduction, or discontinuation of Polivy (see section 4.2).

## Infections

Serious, life threatening or fatal infections, including opportunistic infections, such as pneumonia (including *pneumocystis jirovecii* and other fungal pneumonia), bacteraemia, sepsis, herpes infection, and cytomegalovirus infection have been reported in patients treated with Polivy (see section 4.8). Reactivation of latent infections has been reported. Patients should be closely monitored during treatment for signs of bacterial, fungal, or viral infections and seek medical advice if signs and symptoms appear. Anti-infective prophylaxis should be considered throughout treatment with Polivy. Polivy should not be administered in the presence of an active severe infection. Polivy and any concomitant chemotherapy should be discontinued in patients who develop serious infections.

## Human Immunodeficiency Virus (HIV)

Polivy has not been evaluated in patients with HIV. With regard to co-administration of CYP3A-inhibitors see section 4.5.

## Immunization

Live or live-attenuated vaccines should not be given concurrently with the treatment. Studies have not been conducted in patients who recently received live vaccines.

## Progressive multifocal leukoencephalopathy (PML)

PML has been reported with Polivy treatment (see section 4.8). Patients should be monitored closely for new or worsening neurological, cognitive, or behavioural changes suggestive of PML. Polivy and any concomitant chemotherapy should be withheld if PML is suspected and permanently discontinued if the diagnosis is confirmed.

## Tumour lysis syndrome (TLS)

Patients with high tumour burden and rapidly proliferative tumour may be at increased risk of TLS. Appropriate measures/prophylaxis in accordance with local guidelines should be taken prior to treatment with Polivy. Patients should be monitored closely for TLS during treatment with Polivy.

## Infusion-related reactions

Polivy can cause IRRs, including severe cases. Delayed IRRs as late as 24 hours after receiving Polivy have occurred. An antihistamine and antipyretic should be administered prior to the administration of Polivy, and patients should be monitored closely throughout the infusion. If an IRR occurs, the infusion should be interrupted and appropriate medical management should be instituted (see section 4.2).

## Embryo-foetal toxicity

Based on the mechanism of action and nonclinical studies, Polivy can be harmful to the foetus when administered to a pregnant woman (see section 5.3). Pregnant women should be advised regarding risk to the foetus.

Women of childbearing potential should be advised to use effective contraception during treatment with Polivy and for at least 9 months after the last dose (see section 4.6). Male patients with female partners of childbearing potential should be advised to use effective contraception during treatment with Polivy and for at least 6 months after the last dose (see section 4.6).



## Fertility

In non-clinical studies, polatuzumab vedotin has resulted in testicular toxicity, and may impair male reproductive function and fertility (see section 5.3). Therefore, men being treated with Polivy are advised to have sperm samples preserved and stored before treatment (see section 4.6).

## Elderly

Among 435 previously untreated DLBCL patients treated with Polivy in combination with R-CHP in Study GO39942, 227 (52.2%) were  $\geq 65$  years of age. Patients aged  $\geq 65$  had an incidence of serious adverse reactions of 39.2% and 28.4% in patients aged  $< 65$ . A similar incidence of serious adverse reactions was seen in elderly patients in the R-CHOP treatment arm.

Among 151 previously treated DLBCL patients treated with Polivy in combination with bendamustine and rituximab (BR) in Study GO29365, 103 (68%) were  $\geq 65$  years of age. Patients aged  $\geq 65$  had a similar incidence of serious adverse reactions (55%) to patients aged  $< 65$  (56%). Clinical studies of Polivy did not include sufficient numbers of patients aged  $\geq 65$  to determine whether they respond differently from younger patients.

## Hepatic toxicity

Serious cases of hepatic toxicity that were consistent with hepatocellular injury, including elevations of transaminases and/or bilirubin, have occurred in patients treated with Polivy (see section 4.8). Pre-existing liver disease, elevated baseline liver enzymes, and concomitant medicinal products may increase the risk. Liver enzymes and bilirubin level should be monitored (see section 4.2).

## Excipients

This medicinal product contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

## **4.5 Interaction with other medicinal products and other forms of interaction**

No dedicated clinical drug-drug interaction studies with polatuzumab vedotin in humans have been conducted.

### Drug interactions with concomitant medicines that are CYP3A4 inhibitors, substrates or inducers and co-medications that are P-gp inhibitors

Based on physiological-based pharmacokinetic (PBPK) model simulations of MMAE released from polatuzumab vedotin, strong CYP3A4 and P-gp inhibitors (e.g., ketoconazole) may increase the area under the concentration-time curve (AUC) of unconjugated MMAE by 48%. Caution is advised in case of concomitant treatment with CYP3A4 inhibitor. Patients receiving concomitant strong CYP3A4 inhibitors (e.g., boceprevir, clarithromycin, cobicistat, indinavir, itraconazole, nefazodone, nelfinavir, posaconazole, ritonavir, saquinavir, telaprevir, telithromycin, voriconazole) should be monitored more closely for signs of toxicities.

Unconjugated MMAE is not predicted to alter the AUC of concomitant medicines that are CYP3A4 substrates (e.g., midazolam).

Strong CYP3A4 inducers (e.g., rifampicin, carbamazepine, phenobarbital, phenytoin, St John's wort [*Hypericum perforatum*]) may decrease the exposure of unconjugated MMAE.

## Drug interactions of rituximab, bendamustine, cyclophosphamide, and doxorubicin in combination with polatuzumab vedotin

The pharmacokinetics (PK) of rituximab, bendamustine, cyclophosphamide, and doxorubicin are not affected by co-administration with polatuzumab vedotin. Concomitant rituximab is associated with increased antibody conjugated MMAE (acMMAE) plasma AUC by 24% and decreased unconjugated MMAE plasma AUC by 37%, based on population PK analysis. The plasma AUC of acMMAE and unconjugated MMAE for Polivy plus R-CHP are in line with other studies of Polivy. No dose adjustment is required.

Bendamustine does not affect acMMAE and unconjugated MMAE plasma AUC.

### **4.6 Fertility, pregnancy and lactation**

#### Women of childbearing potential/Contraception in males and females

##### *Women*

Women of childbearing potential should be advised to use effective contraception during treatment with polatuzumab vedotin and for at least 9 months after the last dose.

##### *Men*

Male patients with female partners of childbearing potential should be advised to use effective contraception during treatment with polatuzumab vedotin and for at least 6 months after the last dose.

#### Pregnancy

There are no data in pregnant women using Polivy. Studies in animals have shown reproductive toxicity (see section 5.3). Based on the mechanism of action and nonclinical studies, polatuzumab vedotin can be harmful to the foetus when administered to a pregnant woman. In women of childbearing potential, the pregnancy status shall be checked prior to treatment. Polivy is not recommended during pregnancy and in women of childbearing potential not using contraception unless the potential benefit for the mother outweighs the potential risk to the foetus.

#### Breast-feeding

It is not known whether polatuzumab vedotin or its metabolites are excreted in human breast milk. A risk for breast-feeding children cannot be excluded. Women should discontinue breast-feeding during treatment with Polivy and for at least 3 months after the last dose.

#### Fertility

In nonclinical studies, polatuzumab vedotin has resulted in testicular toxicity, and may impair male reproductive function and fertility (see section 5.3).

Therefore, men being treated with this medicine are advised to have sperm samples preserved and stored before treatment. Men being treated with Polivy are advised not to father a child during treatment and for up to 6 months following the last dose.

### **4.7 Effects on ability to drive and use machines**

Polivy has minor influence on the ability to drive and use machines. IRRs, PN, fatigue, and dizziness may occur during treatment with Polivy (see sections 4.4 and 4.8).

## 4.8 Undesirable effects

### Summary of the safety profile

The safety of Polivy has been evaluated in 435 patients in Study GO39942 (POLARIX). The ADRs described in section 4.8 were identified:

- during treatment and follow-up of previously untreated DLBCL patients from the pivotal clinical trial GO39942 (POLARIX), who received Polivy plus R-CHP (n=435) or R-CHOP (n=438). In the Polivy plus R-CHP group, 91.7% received 6 cycles of Polivy versus 88.5% of patients who received 6 cycles of vincristine in the R-CHOP group.

In previously untreated DLBCL patients treated with Polivy plus R-CHP:

- The most frequently-reported ( $\geq 30\%$ ) adverse drug reactions (ADRs) in patients treated with Polivy plus R-CHP for previously untreated DLBCL were neuropathy peripheral (52.9%), nausea (41.6%), neutropenia (38.4%), and diarrhoea (30.8%).
- Serious adverse reactions were reported in 24.1% of Polivy plus R-CHP treated patients.
- The most common serious adverse reactions reported in  $\geq 5\%$  of patients were febrile neutropenia (10.6%) and pneumonia (5.3%).
- The ADRs leading to treatment regimen discontinuation in  $> 1\%$  of patients treated with Polivy plus R-CHP was pneumonia (1.1%).

The safety of Polivy has been evaluated in 151 patients in Study GO29365. The ADRs described in section 4.8 were identified:

- during treatment and follow-up of previously treated DLBCL patients (n=151) from the pivotal clinical trial GO29365. This includes run-in phase patients (n=6), randomized patients (n=39), and extension cohort patients (n=106) who received Polivy plus BR compared to randomized patients (n=39) who received BR alone. Patients in the treatment arms received a median of 5 cycles of treatment while randomized patients in the comparator arm received a median of 3 cycles of treatment.

In previously treated DLBCL patients treated with Polivy plus BR:

- The most frequently reported ( $\geq 30\%$ ) ADRs (all grades) in patients treated with Polivy plus BR in previously treated DLBCL were neutropenia (45.7%), diarrhoea (35.8%), nausea (33.1%), thrombocytopenia (32.5%), anaemia (31.8%) and neuropathy peripheral (30.5%).
- Serious adverse reactions were reported in 41.7% of Polivy plus BR treated patients.
- The most common serious adverse reactions reported in  $\geq 5\%$  of patients were: febrile neutropenia (10.6%), sepsis (9.9%), pneumonia (8.6%) and pyrexia (7.9%).
- The ADR leading to treatment regimen discontinuation in  $> 5\%$  of patients treated with Polivy plus BR was thrombocytopenia (7.9%).

### Tabulated list of ADRs from clinical trials

The ADRs in 586 patients treated with Polivy are presented in Table 4. The ADRs are listed below by MedDRA system organ class (SOC) and categories of frequency. The corresponding frequency category for each adverse drug reaction is based on the following convention: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1,000$  to  $< 1/100$ ), rare ( $\geq 1/10,000$  to  $< 1/1,000$ ), very rare ( $< 1/10,000$ ). Within each frequency grouping, adverse reactions are presented in the order of decreasing seriousness.

**Table 4** Tabulated list of ADRs in patients treated with Polivy in clinical trials

|   |   |
|---|---|
| <b>Infections and infestations</b>                          |   |
| Very common   | pneumonia <sup>a</sup> , upper respiratory tract infection  |
| Common  | sepsis <sup>a</sup> , herpes virus infection <sup>a</sup> , cytomegalovirus infection, urinary tract infection <sup>c</sup> |
| <b>Blood and lymphatic system disorders</b>                 |   |
| Very common   | febrile neutropenia, neutropenia, thrombocytopenia, anaemia, leukopenia   |
| Common  | lymphopenia, pancytopenia   |
| <b>Metabolism and nutrition disorders</b>                   |   |
| Very common   | hypokalaemia, decreased appetite  |
| Common  | hypocalcaemia, hypoalbuminemia  |
| <b>Nervous system disorders</b>                             |   |
| Very common   | neuropathy peripheral   |
| Common  | dizziness   |
| <b>Eye disorders</b>  |   |
| Uncommon  | vision blurred <sup>b</sup>   |
| <b>Respiratory, thoracic and mediastinal disorders</b>      |   |
| Very common   | cough   |
| Common  | pneumonitis, dyspnoea <sup>c</sup>  |
| <b>Gastrointestinal disorders</b>                           |   |
| Very common   | diarrhoea, nausea, constipation, vomiting, mucositis <sup>c</sup> , abdominal pain  |
| <b>Skin and subcutaneous tissue disorders</b>               |   |
| Very common   | alopecia <sup>c</sup>   |
| Common  | pruritus, skin infections <sup>c</sup> , rash <sup>c</sup> , dry skin <sup>c</sup>  |
| <b>Musculoskeletal disorders</b>                            |   |
| Common  | arthralgia, myalgia <sup>c</sup>  |
| <b>General disorders and administration site conditions</b> |   |
| Very common   | pyrexia, fatigue, asthenia  |
| Common  | peripheral edema <sup>c</sup> , chills  |
| <b>Investigations</b>                                       |   |
| Very common   | weight decreased  |
| Common  | transaminases increased, lipase increase <sup>b</sup> , hypophosphataemia   |

| <b>Injury, poisoning and procedural complications</b> |                           |
|---|---------------------------|
| Very Common   | infusion related reaction |

<sup>a</sup> ADR associated with fatal outcome

<sup>b</sup> ADRs observed in relapsed or refractory DLBCL only.

<sup>c</sup> ADRs observed in previously untreated DLBCL only.

The listed ADRs were observed in both previously untreated DLBCL and relapsed or refractory DLBCL except where indicated with footnotes.

Rare and very rare ADRs: none

### Description of selected adverse drug reactions

#### *Myelosuppression*

In a placebo-controlled study GO39942 (POLARIX), 0.5% of patients in the Polivy plus R-CHP arm discontinued study treatment due to neutropenia. No patients discontinued study treatment in the R-CHOP arm due to neutropenia. Thrombocytopenia events led to discontinuation of study treatment in 0.2% of patients in the Polivy plus R-CHP arm compared to no patients in the R-CHOP arm. No patients discontinued treatment due to anaemia in either the Polivy plus R-CHP arm or R-CHOP arm.

In an open-label study GO29365, 4% of patients in the Polivy plus BR arms discontinued Polivy due to neutropenia compared to 2.6% of patients in the BR arm who discontinued treatment due to neutropenia. Thrombocytopenia events led to discontinuation of treatment in 7.9% of patients in the Polivy plus BR arms and 5.1% of patients in the BR arm. No patients discontinued treatment due to anaemia in either the Polivy plus BR arms or BR arm. In the Polivy plus BR arms, Grade 3 or higher neutropenia, thrombocytopenia, and anaemia were reported in 40.4%, 25.8%, and 12.6% of patients, respectively.

#### *Peripheral neuropathy (PN)*

In a placebo-controlled study GO39942 (POLARIX), in the Polivy plus R-CHP arm, Grade 1, 2, and 3 PN were reported in 39.1%, 12.2% and 1.6% of patients, respectively. In the R-CHOP arm, Grade 1, 2 and 3 PN events were reported in 37.2%, 15.5% and 1.1% of patients, respectively. No Grade 4-5 PN events were reported in either the Polivy plus R-CHP arm or R-CHOP arm. 0.7% of patients discontinued study treatment in the Polivy plus R-CHP arm due to PN compared to 2.3% in the R-CHOP arm. 4.6% of patients in the Polivy plus R-CHP arm had study treatment dose reduction due to PN compared to 8.2% in the R-CHOP arm. In the Polivy plus R-CHP arm, the median time to onset of first event of PN was 2.27 months compared to 1.87 months in the R-CHOP arm. PN events resolved in 57.8% of patients in the Polivy plus R-CHP arm as of the clinical cut off date compared to 66.9% in the R-CHOP arm. The median time to peripheral neuropathy resolution was 4.04 months in the Polivy plus R-CHP arm compared to 4.6 months in the R-CHOP arm.

In an open-label study GO29365, in the Polivy plus BR arms, Grade 1 PN and Grade 2 PN were reported in 15.9% and 12.6% of patients, respectively. In the BR arm, Grade 1 and 2 PN events were reported in 2.6% and 5.1% of patients, respectively. One Grade 3 PN event was reported in the Polivy plus BR arms and no patients reported PN events in the BR arm. No Grade 4-5 PN events were reported in either the Polivy plus BR arms or BR arm. 2.6% of patients discontinued Polivy treatment due to PN and 2.0% of patients had Polivy dose reduction due to PN. No patients in the BR arm discontinued treatment or had dose reductions due to PN. In the Polivy plus BR arms, the median time to onset of first event of PN was 1.6 months, and 39.1% of patients with PN events reported event resolution.

#### *Infections*

In a placebo-controlled study GO39942 (POLARIX), infections, including pneumonia and other types of infections, were reported in 49.7% of patients in the Polivy plus R-CHP arm and 42.7% of patients in the R-CHOP arm. Grade 3-4 infections occurred in 14.0% of patients in the Polivy plus R-CHP arm and 11.2% of patients in the R-CHOP arm. In the Polivy plus R-CHP arm, serious infections were reported in 14.0% of patients and fatal infections were reported in 1.1% of patients. In the R-CHOP arm, serious infections were reported in 10.3% of patients and fatal infections were reported in 1.4%

of patients. 7 patients (1.6%) in the Polivy plus R-CHP arm discontinued treatment due to infection compared to 10 patients (2.3%) in the R-CHOP arm.

In an open-label study GO29365, infections, including pneumonia and other types of infections, were reported in 48.3% of patients in the Polivy plus BR arms and 51.3% of patients in the BR arm. In the Polivy plus BR arms, serious infections were reported in 27.2% of patients and fatal infections were reported in 6.6% of patients. In the BR arm, serious infections were reported in 30.8% of patients and fatal infections were reported in 10.3% of patients. Four patients (2.6%) in the Polivy plus BR arms discontinued treatment due to infection compared to 2 patients (5.1%) in the BR arm.

#### *Progressive multifocal leukoencephalopathy (PML)*

In a placebo-controlled study GO39942 (POLARIX), no cases of PML were reported.

In an open-label study GO29365, one case of PML, which was fatal, occurred in one patient treated with Polivy plus bendamustine and obinutuzumab. This patient had three prior lines of therapy that included anti-CD20 antibodies.

#### *Hepatic toxicity*

In a placebo-controlled study GO39942 (POLARIX), hepatic toxicity was reported in 10.6% of patients in the Polivy plus R-CHP arm and 7.3% of patients in the R-CHOP arm. In the Polivy plus R-CHP arm, most events were Grade 1–2 (8.7%); Grade 3 events were reported in 1.8% of patients. There were no Grade 4 or 5 events. Serious hepatic toxicity events were reported in 1 patient (0.2%) and were reversible.

In another study, two cases of serious hepatic toxicity (hepatocellular injury and hepatic steatosis) were reported and were reversible.

#### *Gastrointestinal toxicity*

In a placebo-controlled study GO39942 (POLARIX), gastrointestinal toxicity events were reported in 76.1% of patients in the Polivy plus R-CHP arm compared to 71.9% of patients in the R-CHOP arm. Most events were Grade 1–2, and Grade  $\geq 3$  events were reported in 9.7% of patients in the Polivy plus R-CHP arm compared to 8.2% of patients in the R-CHOP arm. The most common gastrointestinal toxicity events were nausea and diarrhoea.

In an open-label study GO29365, gastrointestinal toxicity events were reported in 72.8% of patients in the Polivy plus BR arms compared to 66.7% of patients in the BR arm. Most events were Grade 1-2, and Grade 3-4 events were reported in 16.5% of patients in the Polivy plus BR arms compared to 12.9% of patients in the BR arm. The most common gastrointestinal toxicity events were diarrhoea and nausea.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via **the national reporting system** listed in [Appendix V](#).

#### **4.9 Overdose**

There is no experience with overdose in human clinical trials. The highest dose tested to date is 2.4 mg/kg administered as an intravenous infusion; it was associated with a higher frequency and severity of PN events. Patients who experience overdose should have immediate interruption of their infusion and be closely monitored.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: antineoplastic agents; other antineoplastic agents; monoclonal antibodies  
ATC code: L01FX14

#### Mechanism of action

Polatuzumab vedotin is a CD79b-targeted antibody-drug conjugate that preferentially delivers a potent anti-mitotic agent (monomethyl auristatin E, or MMAE) to B-cells, which results in the killing of malignant B-cells. The polatuzumab vedotin molecule consists of MMAE covalently attached to a humanized immunoglobulin G1 monoclonal antibody via a cleavable linker. The monoclonal antibody binds with high affinity and selectivity to CD79b, a cell surface component of the B-cell receptor. CD79b expression is restricted to normal cells within the B-cell lineage (with the exception of plasma cells) and malignant B-cells; it is expressed in > 95% of diffuse large B-cell lymphoma. Upon binding CD79b, polatuzumab vedotin is rapidly internalized and the linker is cleaved by lysosomal proteases to enable intracellular delivery of MMAE. MMAE binds to microtubules and kills dividing cells by inhibiting cell division and inducing apoptosis.

#### Pharmacodynamic effects

##### *Cardiac electrophysiology*

Polatuzumab vedotin did not prolong the mean QTc interval to any clinically relevant extent based on ECG data from two open-label studies in patients with previously treated B-cell malignancies at the recommended dosage.

#### Clinical efficacy and safety

##### *Previously untreated DLBCL*

The efficacy of Polivy was evaluated in an international, multicenter, randomized double-blind, placebo-controlled study (POLARIX, GO39942) in 879 patients with previously untreated DLBCL.

Eligible patients were age 18–80, and had IPI score 2–5, and ECOG Performance Status 0–2. Histologies included DLBCL (not otherwise specified (NOS), activated B-cell (ABC), germinal center B-cell (GCB)), HGBL (NOS, double-hit, triple-hit), and other large B-cell lymphoma subtypes (EBV positive, T-cell rich/histiocyte rich). Patients did not have known CNS lymphoma or peripheral neuropathy > Grade 1.

Patients were randomized 1:1 to receive Polivy plus R-CHP or R-CHOP for six 21-day cycles followed by two additional cycles of rituximab alone in both arms. Patients were stratified by IPI score (2 vs 3-5), presence or absence of bulky disease (lesion  $\geq$  7.5 cm), and geographical region.

Polivy was administered intravenously at 1.8 mg/kg on Day 1 of Cycles 1–6. R-CHP or R-CHOP were administered starting on Day 1 of Cycles 1–6 followed by rituximab alone on Day 1 of Cycles 7–8.

Dosing in each treatment arm was administered according to the following:

- Polivy + R-CHP arm: Polivy 1.8 mg/kg, rituximab 375 mg/m<sup>2</sup>, cyclophosphamide 750 mg/m<sup>2</sup>, doxorubicin 50 mg/m<sup>2</sup>, and prednisone 100 mg/day, on Days 1-5 of every cycle, orally.
- R-CHOP arm: rituximab 375 mg/m<sup>2</sup>, cyclophosphamide 750 mg/m<sup>2</sup>, doxorubicin 50 mg/m<sup>2</sup>, vincristine 1.4 mg/m<sup>2</sup>, and prednisone 100 mg/day, on Days 1-5 of every cycle, orally.

The two treatment groups were generally balanced with respect to baseline demographics and disease characteristics. The median age was 65 years (range 19 to 80 years), 53.6% of patients were white and 53.8% were male, 43.8% had bulky disease, 38.0% had IPI score 2, 62.0% had IPI score 3–5, and 88.7% had Stage 3 or 4 disease. The majority of patients (84.2%) had DLBCL (including NOS, ABC, and GCB).

211 patients did not have a cell of origin (COO) result reported. Of the COO evaluable population (n=668), 33.1% of patients had ABC like DLBCL and 52.7% of patients had GCB like DLBCL, by gene expression profiling.

The primary endpoint of the study was investigator-assessed progression free survival. The median duration of follow up was 28.2 months. Efficacy results are summarized in Table 5 and in Figure 1.

**Table 5 Summary of efficacy in patients with previously untreated DLBCL from Study GO39942 (POLARIX)**

|  | <b>Polivy + R-CHP<br/>N=440</b> | <b>R-CHOP<br/>N=439</b> |
|--|---------------------------------|-------------------------|
| <b>Primary Endpoint</b>  |                                 |                         |
| Progression free survival <sup>1,*</sup>                       |                                 |                         |
| Number (%) of patients with events                             | 107 (24.3%)                     | 134 (30.5%)             |
| HR (95% CI)  | 0.73 [0.57, 0.95]               |                         |
| p-value <sup>3,**</sup>  | 0.0177                          |                         |
| 2-year PFS estimate (%)  | 76.7                            | 70.2                    |
| [95% CI]   | [72.65, 80.76]                  | [65.80, 74.61]          |
| <b>Key secondary endpoints</b>                                 |                                 |                         |
| Event-free survival (EFS <sub>eff</sub> ) <sup>1</sup>         |                                 |                         |
| Number (%) of patients with event                              | 112 (25.5%)                     | 138 (31.4%)             |
| HR [95% CI]  | 0.75 [0.58, 0.96]               |                         |
| p-value <sup>3,**</sup>  | 0.0244                          |                         |
| Objective Response Rate (ORR) at End of Treatment <sup>2</sup> |                                 |                         |
| Responders (%) (CR, PR)  | 376 (85.5%)                     | 368 (83.8%)             |
| Difference in response rate (%) [95% CI]                       | 1.63 [-3.32, 6.57]              |                         |
| Complete Response (%) (CR) Rate <sup>2,*</sup>                 |                                 |                         |
| Responders (%)   | 343 (78.0%)                     | 325 (74.0%)             |
| Difference in response rate (%) [95% CI]                       | 3.92 [-1.89, 9.70]              |                         |
| Partial response (%) (PR)                                      | 33 (7.5%)                       | 43 (9.8%)               |
| 95% CI Clopper-Pearson   | [5.22, 10.37]                   | [7.18, 12.97]           |

INV: Investigator; BICR: Blinded independent central review; CI: Confidence interval; HR: Hazard ratio; PFS: Progression free survival; EFS<sub>eff</sub>: Event free survival efficacy: used to reflect EFS events that are due to efficacy and defined as time from date of randomization to the earliest occurrence of any of the following: disease progression/relapse, death due to any cause, the primary efficacy reason determined by the investigator, other than disease progression/relapse, that led to initiation of any non-protocol specified anti-lymphoma treatment (NALT), if biopsy was obtained after treatment completion and was positive for residual disease regardless of whether NALT was initiated or not; CMH: Cochran-Mantel-Haenszel

1) INV-assessed

2) BICR-assessed

3) Log-rank test, stratified


\*Per Lugano 2014 Response Criteria

\*\*Stratified by IPI (2 vs 3-5), presence or absence of bulky disease, geography

At the interim analysis, the key secondary endpoint of OS was immature and was not statistically different [stratified hazard ratio of 0.94 (95% CI, 0.65, 1.37); p=0.7524].



**Figure 1 Kaplan Meier curve of INV-assessed progression-free survival (PFS) in Study GO39942 (POLARIX)**



#### *Relapsed or refractory DLBCL*

The efficacy of Polivy was evaluated in an international, multicentre, open-label study (GO29365) which included a randomized cohort of 80 patients with previously treated DLBCL. Patients were randomized 1:1 to receive Polivy plus BR or BR alone for six 21-day cycles. Patients were stratified by duration of response to last prior treatment of  $\leq 12$  months or  $> 12$  months.

Eligible patients were not candidates for autologous haematopoietic stem cell transplant (HSCT) and had relapsed or refractory disease after receiving at least one prior systemic chemotherapy regimen. The study excluded patients with prior allogeneic HSCT, central nervous system lymphoma, transformed indolent lymphoma, grade 3b FL, significant cardiovascular or pulmonary disease, active infections, AST or alanine transaminase (ALT)  $> 2.5 \times$  ULN or total bilirubin  $\geq 1.5 \times$  ULN, creatinine  $> 1.5 \times$  ULN (or CrCl  $< 40$  mL/min) unless due to underlying lymphoma.

Polivy was given intravenously at 1.8 mg/kg administered on Day 2 of Cycle 1 and on Day 1 of Cycles 2-6. Bendamustine was administered at 90 mg/m<sup>2</sup> intravenously daily on Days 2 and 3 of Cycle 1 and on Days 1 and 2 of Cycles 2-6. Rituximab was administered at 375 mg/m<sup>2</sup> on Day 1 of Cycles 1-6.

Among the 80 patients who were randomized to receive Polivy plus BR (n=40) or BR alone (n=40) the majority were white (71%) and male (66%). The median age was 69 years (range: 30-86 years). Sixty-four out of 80 patients (80%) had ECOG performance status (PS) of 0-1 and 14 out of 80 patients (18%) had ECOG PS of 2. The majority of patients (98%) had DLBCL not otherwise specified (NOS). Overall, 48% of patients had activated B-cell (ABC) DLBCL and 40% had germinal center B-cell like (GCB) DLBCL. Primary reasons patients were not candidates for HSCT included age (40%), insufficient response to salvage therapy (26%) and prior transplant failure (20%). The median number of prior therapies was 2 (range: 1-7), with 29% (n=23) receiving one prior therapy, 25% (n=20) receiving 2 prior therapies, and 46% (n=37) receiving 3 or more prior therapies. All except one patient in the pola+BR arm of the randomized Phase II were naïve to bendamustine treatment. 80% of patients had refractory disease. For patients who received polatuzumab vedotin plus BR and had CD3+ lymphocyte count evaluated, the absolute CD3+ lymphocyte count was

> 200 cells/ $\mu$ L in 95%, 79% and 83% of patients analyzed at prior to therapy (n=134), end of treatment (n=72) and 6 months after end of treatment (n=18), respectively.

The primary endpoint of the study was complete response (CR) rate at end of treatment (6-8 weeks after Day 1 of Cycle 6 or last study treatment) as assessed by PET-CT by an Independent Review Committee (IRC).

**Table 6 Summary of efficacy in patients with previously treated DLBCL from study GO29365**

|  | <b>Polivy + bendamustine<br/>+ rituximab<br/>N=40</b> | <b>Bendamustine +<br/>rituximab<br/>N=40</b> |
|--|---|--|
|  | <b>Median observation time 22 months</b>              |  |
| <b>Primary endpoint</b>                                      |   |  |
| Complete Response Rate* (IRC-assessed) at End of treatment** |   |  |
| Responders (%)   | 16 (40.0)   | 7 (17.5)                                     |
| Difference in response rate (%) [95% CI]                     | 22.5 [2.6, 40.2]                                      |  |
| p-value (CMH chi-squared test***)                            | 0.0261  |  |
| <b>Key secondary and exploratory endpoints</b>               |   |  |
| Duration of response (INV-assessed)                          |   |  |
| Number of patients included in analysis                      | 28  | 13   |
| Number (%) of patients with event                            | 17 (60.7)   | 11 (84.6)                                    |
| Median DOR (95% CI), months                                  | 10.3 (5.6, NE)  | 4.1 (2.6, 12.7)                              |
| HR [95% CI]  | 0.44 [0.20, 0.95]                                     |  |
| p-value (Log-Rank test, stratified***)                       | 0.0321  |  |
| Overall Response Rate* (INV-assessed) at End of Treatment**  |   |  |
| Responders (%) (CR, PR)                                      | 19 (47.5)   | 7 (17.5)                                     |
| Difference in response rate (%) [95% CI]                     | 30.0 [9.5, 47.4]                                      |  |
| p-value (CMH chi-squared test***)                            | 0.0036  |  |
| Complete Response (%) (CR)                                   | 17 (42.5)   | 6 (15.0)                                     |
| Difference in response rate (%) [95% CI]                     | 27.5 [7.7, 44.7]                                      |  |
| p-value (CMH chi-squared test***)                            | 0.0061  |  |
| Partial Response (%) (PR)                                    | 2 (5.0)   | 1 (2.5)                                      |
| 95% CI Clopper-Pearson                                       | [0.6, 16.9]   | [0.06, 13.2]                                 |
| Best Overall Response Rate* (INV-assessed)                   |   |  |
| Responders (%) (CR, PR)                                      | 28 (70.0)   | 13 (32.5)                                    |
| Difference in response rate (%) [95% CI]                     | 37.5 [15.6, 54.7]                                     |  |
| Complete Response (%) (CR)                                   | 23 (57.5)   | 8 (20.0)                                     |
| 95% CI Clopper-Pearson                                       | [40.9, 73.0]  | [9.1, 35.7]                                  |
| Partial Response (%) (PR)                                    | 5 (12.5)  | 5 (12.5)                                     |
| 95% CI Clopper-Pearson                                       | [4.2, 26.8]   | [4.2, 26.8]                                  |

CI: Confidence Interval; CMH: Cochran-Mantel-Haenszel; CR: Complete Response; DOR: Duration of Response; HR: Hazard Ratio; INV: Investigator; IRC: Independent Review Committee; NE: Not evaluable; PR: Partial Response

\*Per modified Lugano 2014 criteria: Bone marrow confirmation of PET-CT CR required. PET-CT PR required meeting both PET-CT criteria and CT criteria.

\*\*6-8 weeks after Day 1 of Cycle 6 or last study treatment

\*\*\* Stratification by duration of response to prior therapy ( $\leq$  12 months vs  $>$  12 months)

Overall survival (OS) was an exploratory endpoint which was not type 1 error controlled. The median OS in the Polivy plus BR arm was 12.4 months (95% CI: 9.0, NE) vs 4.7 months (95% CI: 3.7, 8.3) in

the control arm. The unadjusted estimate for OS HR was 0.42. When accounting for the influence of baseline covariates the OS HR was adjusted to 0.59. Covariates included primary refractory status, number of prior lines of therapy, IPI, and prior stem cell transplant.

Investigator-assessed progression free survival (PFS) was an exploratory endpoint which was not type 1 error controlled. The median PFS in the Polivy plus BR arm was 7.6 months (95% CI: 6.0, 17.0) vs 2.0 months (95% CI: 1.5, 3.7) in the control arm. The unadjusted estimate for PFS HR was 0.34.

### Immunogenicity

As with all therapeutic proteins, there is the potential for an immune response in patients treated with polatuzumab vedotin. In Studies GO39442 (POLARIX) and GO29365, 1.4% (6/427) and 5.2% (12/233) of patients tested positive for antibodies against polatuzumab vedotin, respectively, of which none were positive for neutralizing antibodies. Due to the limited number of anti-polatuzumab vedotin antibody positive patients, no conclusions can be drawn concerning a potential effect of immunogenicity on efficacy or safety.

Immunogenicity assay results are highly dependent on several factors including assay sensitivity and specificity, assay methodology, sample handling, timing of sample collection, concomitant medications and underlying disease. For these reasons, comparison of incidence of antibodies to polatuzumab vedotin with the incidence of antibodies to other products may be misleading.

### Paediatric population

The European Medicines Agency has waived the obligation to submit results of studies with Polivy in all subsets of the paediatric population for the treatment of mature B-cell neoplasms (see section 4.2 for information on paediatric use).

## **5.2 Pharmacokinetic properties**

Antibody-conjugated MMAE (acMMAE) plasma exposure increased dose-proportionally over the 0.1 to 2.4 mg/kg polatuzumab vedotin dose range. After the first 1.8 mg/kg polatuzumab vedotin dose, the acMMAE mean maximum concentration ( $C_{max}$ ) was 803 ( $\pm 233$ ) ng/mL and the area under the concentration-time curve from time zero to infinity ( $AUC_{inf}$ ) was 1860 ( $\pm 966$ ) day•ng/mL. Based on the population PK analysis, Cycle 3 acMMAE AUC increased by approximately 30% over Cycle 1 AUC, and achieved more than 90% of the Cycle 6 AUC. The terminal half-life at Cycle 6 was approximately 12 days (95% CI of 8.1-19.5 days) for acMMAE. Based on population PK analysis, the predicted acMMAE concentration at the end of Cycle 6 is approximately 80% of the theoretical steady-state value. Exposures of unconjugated MMAE, the cytotoxic component of polatuzumab vedotin, increased dose proportionally over the 0.1 to 2.4 mg/kg polatuzumab vedotin dose range. MMAE plasma concentrations followed formation rate limited kinetics. After the first 1.8 mg/kg polatuzumab vedotin dose, the  $C_{max}$  was 6.82 ( $\pm 4.73$ ) ng/mL, the time to maximum plasma concentration is approximately 2.5 days, and the terminal half-life is approximately 4 days. Plasma exposures of unconjugated MMAE are < 3% of acMMAE exposures. Based on the population PK analysis there is a decrease of plasma unconjugated MMAE exposure (AUC) after repeated every-three-week dosing.

Based on population pharmacokinetics simulations, a post-hoc analysis predicted exposure to unconjugated MMAE for patients with bodyweight over 100 kg to be increased by not more than 55%.

### Absorption

Polivy is administered as an intravenous infusion. There have been no studies performed with other routes of administration.

## Distribution

The population estimate of central volume of distribution for acMMAE was 3.15 L, which approximated plasma volume. *In vitro*, MMAE is moderately bound (71%-77%) to human plasma proteins. MMAE does not significantly partition into human red blood cells *in vitro*; the blood to plasma ratio is 0.79 to 0.98.

*In vitro* data indicate that MMAE is a P-gp substrate but does not inhibit P-gp at clinically relevant concentrations.

## Biotransformation

Polatuzumab vedotin is expected to undergo catabolism in patients, resulting in the production of small peptides, amino acids, unconjugated MMAE, and unconjugated MMAE related catabolites. The levels of MMAE metabolites have not been measured in human plasma.

*In vitro* studies indicate that MMAE is a substrate for CYP3A4/5 but does not induce major CYP enzymes. MMAE is a weak time-dependent inhibitor of CYP3A4/5 but does not competitively inhibit CYP3A4/5 at clinically relevant concentrations.

MMAE does not inhibit CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, or CYP2D6.

## Elimination

Based on a population PK analysis, the conjugate (acMMAE) is primarily eliminated by non-specific linear clearance pathway with a value of 0.9 L/day. *In vivo* studies in rats dosed with polatuzumab vedotin (radiolabel on MMAE) demonstrate that the majority of radioactivity is excreted in faeces and the minority of radioactivity is excreted in urine.

## Paediatric population

No studies have been conducted to investigate the pharmacokinetics of polatuzumab vedotin in the paediatric population (<18 years old).

## Elderly

Age did not have an effect on the pharmacokinetics of acMMAE and unconjugated MMAE based on population PK analyses with patients aged 19-89 years. No significant difference was observed in the pharmacokinetics of acMMAE and unconjugated MMAE among patients < 65 years of age (n=394) and patients ≥ 65 years of age (n=495) based on population PK analyses.

## Renal impairment

In patients with mild (CrCL 60-89 mL/min, n=361) or moderate (CrCL 30-59 mL/min, n=163) renal impairment, acMMAE and unconjugated MMAE exposures are similar to patients with normal renal function (CrCL ≥ 90 mL/min, n=356), based on population PK analyses. There are insufficient data to assess the impact of severe renal impairment (CrCL 15-29 mL/min, n=4) on PK. No data are available in patients with end-stage renal disease and/or who are on dialysis.

## Hepatic impairment

In patients with mild hepatic impairment [AST or ALT > 1.0 to 2.5 × ULN or total bilirubin > 1.0 to 1.5 × ULN, n=133], acMMAE exposures are similar whereas unconjugated MMAE AUC are not more than 40% higher compared to patients with normal hepatic function (n=737), based on population PK analyses.

There are insufficient data to assess the impact of moderate hepatic impairment (total bilirubin  $> 1.5\text{-}3 \times \text{ULN}$ ,  $n=11$ ) on PK. Limited data are available in patients with severe hepatic impairment or liver transplantation.

### **5.3 Preclinical safety data**

#### Systemic toxicity

In both rats and cynomolgus monkeys, the predominant systemic toxicities associated with administration of MMAE and polatuzumab vedotin included reversible bone marrow toxicity and associated peripheral blood cell effects.

#### Genotoxicity

No dedicated mutagenicity studies have been performed with polatuzumab vedotin. MMAE was not mutagenic in the bacterial reverse mutation assay (Ames test) or the L5178Y mouse lymphoma forward mutation assay.

MMAE was genotoxic in the rat bone marrow micronucleus study probably through an aneugenic mechanism. This mechanism is consistent with the pharmacological effect of MMAE as a microtubule disrupting agent.

#### Carcinogenicity

No dedicated carcinogenicity studies have been performed with polatuzumab vedotin and/or MMAE.

#### Impairment of fertility

No dedicated fertility studies in animals have been performed with polatuzumab vedotin. However, results of the 4-week rat toxicity study indicate the potential for polatuzumab vedotin to impair male reproductive function and fertility. Testicular seminiferous tubule degeneration did not reverse following a 6-week treatment-free period and correlated with decreased testes weight and gross findings at recovery necropsy of small and/or soft testes in males given  $\geq 2$  mg/kg.

#### Reproductive toxicity

No dedicated teratogenicity studies in animals have been performed with polatuzumab vedotin. However, treatment of pregnant rats with MMAE at 0.2 mg/kg caused embryoletality and foetal malformations (including protruding tongue, malrotated limbs, gastroschisis, and agnathia). Systemic exposure (AUC) in rats at a dose of 0.2 mg/kg MMAE is approximately 50% of the AUC in patients who received the recommended dose of 1.8 mg/kg Polivy every 21 days.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Succinic acid  
Sodium hydroxide (for pH-adjustment)  
Sucrose  
Polysorbate 20 (E 432)

### **6.2 Incompatibilities**

This medicinal product must not be mixed or diluted with other medicinal products except those mentioned in section 6.6.

### 6.3 Shelf life

#### Unopened vial

30 months

#### Reconstituted solution

From a microbiological point of view, the reconstituted solution should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours refrigerated (2 °C-8 °C), unless reconstitution has taken place in controlled and validated aseptic conditions. Chemical and physical in-use stability of the reconstituted solution has been demonstrated for up to 72 hours refrigerated (2 °C-8 °C) and up to 24 hours at room temperature (9 °C-25 °C).

#### Diluted solution

From a microbiological point of view, the prepared solution for infusion should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours refrigerated (2 °C-8 °C), unless dilution has taken place in controlled and validated aseptic conditions. Chemical and physical stability of the prepared solution for infusion has been demonstrated for the durations listed in Table 7. The diluted solution must be discarded if storage time exceeds the limits specified in Table 7.

**Table 7 Durations for which chemical and physical stability of the prepared solution for infusion have been demonstrated**

| <b>Diluent used to prepare solution for infusion</b> | <b>Solution for infusion storage conditions<sup>1</sup></b>                                   |
|--|---|
| Sodium chloride 9 mg/mL (0.9%)                       | Up to 72 hours refrigerated (2 °C – 8 °C) or up to 4 hours at room temperature (9 °C – 25 °C) |
| Sodium chloride 4.5 mg/mL (0.45%)                    | Up to 72 hours refrigerated (2 °C – 8 °C) or up to 8 hours at room temperature (9 °C – 25 °C) |
| 5% glucose   | Up to 72 hours refrigerated (2 °C – 8 °C) or up to 8 hours at room temperature (9 °C – 25 °C) |

<sup>1</sup> To ensure product stability, do not exceed specified storage durations.

### 6.4 Special precautions for storage

Store in a refrigerator (2 °C- 8 °C).

Do not freeze.

Keep the vial in the outer carton in order to protect from light.

For storage conditions after reconstitution and dilution of the medicinal product, see section 6.3.

### 6.5 Nature and contents of container

#### Polivy 30 mg powder for concentrate for solution for infusion:

6 mL vial (colourless Type 1 glass) closed with a stopper (fluororesin laminate), with an aluminium seal with plastic flip-off cap containing 30 mg polatuzumab vedotin. Pack size of one vial.

### Polivy 140 mg powder for concentrate for solution for infusion:

20 mL vial (colourless Type 1 glass) closed with a stopper (fluororesin laminate), with an aluminium seal with plastic flip-off cap containing 140 mg polatuzumab vedotin. Pack size of one vial.

## **6.6 Special precautions for disposal and other handling**

### General precautions

Polivy contains a cytotoxic component. To be administered under the supervision of a physician experienced in the use of cytotoxic agents. Procedures for proper handling and disposal of antineoplastic and cytotoxic medicines should be used.

The reconstituted product contains no preservative and is intended for single-dose only. Proper aseptic technique throughout the handling of this medicinal product should be followed.

Polivy must be reconstituted using sterile water for injection and diluted into an intravenous infusion bag containing sodium chloride 9 mg/mL (0.9%) solution for injection, or sodium chloride 4.5 mg/ml (0.45%) solution for injection, or 5% glucose prior to administration.

The reconstituted solution and solution for infusion should not be frozen or exposed to direct sunlight.

### Instructions for reconstitution

- Polivy 30 mg: Using a sterile syringe, slowly inject 1.8 mL of sterile water for injection into the 30 mg Polivy vial to yield a single-dose solution containing 20 mg/mL polatuzumab vedotin. Direct the stream toward the wall of the vial and not directly on the lyophilized cake.
- Polivy 140 mg: Using a sterile syringe, slowly inject 7.2 mL of sterile water for injection into the 140 mg Polivy vial to yield a single-dose solution containing 20 mg/mL polatuzumab vedotin. Direct the stream toward the wall of the vial and not directly on the lyophilized cake.
- Swirl the vial gently until completely dissolved. Do not shake.
- Inspect the reconstituted solution for discoloration and particulate matter. The reconstituted solution should appear colourless to slightly brown, clear to slightly opalescent, and free of visible particulates. Do not use if the reconstituted solution is discoloured, is cloudy, or contains visible particulates.

### Instructions for dilution

1. Polivy must be diluted to a final concentration of 0.72-2.7 mg/mL in an intravenous infusion bag, with a minimum volume of 50 mL, containing 9 mg/mL sodium chloride solution for injection, or 4.5 mg/mL sodium chloride solution for injection, or 5% glucose.
2. Determine the volume of 20 mg/mL reconstituted solution needed based on the required dose (see below):

$$\text{Total Polivy dose (mL) to be further diluted} = \frac{\text{Polivy dose (mg/kg)} \times \text{patient's weight (kg)}}{\text{Reconstituted vial concentration (20 mg/mL)}}$$

3. Withdraw the required volume of reconstituted solution from the Polivy vial using a sterile syringe and dilute into the intravenous infusion bag. Discard any unused portion left in the vial.
4. Gently mix the intravenous bag by slowly inverting the bag. Do not shake.
5. Inspect the intravenous bag for particulates and discard if present.

Avoid transportation of the prepared solution for infusion as agitation stress can result in aggregation. If the prepared infusion will be transported, remove air from the infusion bag and limit transportation to 30 minutes room temperature (9°C – 25°C) or 24 hours refrigerated (2°C – 8°C). If air is removed, an infusion set with a vented spike is required to ensure accurate dosing during the infusion. The total storage plus transportation times of the diluted product should not exceed the storage duration specified in Table 7 (see section 6.3).

Polivy must be administered using a dedicated infusion line equipped with sterile, non-pyrogenic, low-protein binding in-line or add-on filter (0.2 or 0.22 micrometer pore size) and catheter.

Polivy is compatible with intravenous infusion bags with product contacting materials of polyvinyl chloride (PVC) or polyolefins such as polyethylene (PE) and polypropylene. In addition, no incompatibilities have been observed with infusion sets or infusion aids with product contacting materials of PVC, PE, polyurethane, polybutadiene, acrylonitrile butadiene styrene, polycarbonate, polyetherurethane, fluorinated ethylene propylene, or polytetrafluorethylene and with filter membranes composed of polyether sulfone or polysulfone.

#### Disposal

Polivy is for single-use only.

Any unused product or waste material should be disposed of in accordance with local requirements.

### **7. MARKETING AUTHORISATION HOLDER**

Roche Registration GmbH  
Emil-Barell-Strasse 1  
79639 Grenzach-Wyhlen  
Germany

### **8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/19/1388/001  
EU/1/19/1388/002

### **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 16 January 2020  
Date of latest renewal: 03 December 2021

### **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.



## **ANNEX II**

- A. MANUFACTURER(S) OF THE BIOLOGICAL ACTIVE SUBSTANCE(S) AND MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

**A. MANUFACTURER OF THE BIOLOGICAL ACTIVE SUBSTANCE AND MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturer of the biological active substance

Lonza Ltd.  
Lonzastrasse  
CH-3930 Visp  
Switzerland

Name and address of the manufacturer responsible for batch release

Roche Pharma AG  
Emil-Barell-Strasse 1  
79639 Grenzach-Wyhlen  
Germany

**B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

**C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

- **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

**D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

- **Risk management plan (RMP)**

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Polivy 140 mg powder for concentrate for solution for infusion  
polatuzumab vedotin

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each vial contains 140 mg of polatuzumab vedotin.  
After reconstitution each mL contains 20 mg of polatuzumab vedotin.

**3. LIST OF EXCIPIENTS**

succinic acid, sodium hydroxide, sucrose, polysorbate 20.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Powder for concentrate for solution for infusion  
1 vial

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

For intravenous use after reconstitution and dilution  
Read the package leaflet before use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

Cytotoxic  
Do not shake

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator  
Do not freeze  
Keep the vial in the outer carton in order to protect from light

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Roche Registration GmbH  
Emil-Barell-Strasse 1  
79639 Grenzach-Wyhlen  
Germany

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/19/1388/001

**13. BATCH NUMBER**

Batch

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Justification for not including Braille accepted

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**VIAL**

**1. NAME OF THE MEDICINAL PRODUCT AND ROUTE OF ADMINISTRATION**

Polivy 140 mg powder for concentrate  
polatuzumab vedotin  
Intravenous use

**2. METHOD OF ADMINISTRATION**

IV after reconstitution and dilution

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

140 mg

**6. OTHER**

Cytotoxic

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Polivy 30 mg powder for concentrate for solution for infusion  
polatuzumab vedotin

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each vial contains 30 mg of polatuzumab vedotin.  
After reconstitution each mL contains 20 mg of polatuzumab vedotin.

**3. LIST OF EXCIPIENTS**

succinic acid, sodium hydroxide, sucrose, polysorbate 20.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Powder for concentrate for solution for infusion  
1 vial

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

For intravenous use after reconstitution and dilution  
Read the package leaflet before use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

Cytotoxic  
Do not shake

**8. EXPIRY DATE**

EXP



**9. SPECIAL STORAGE CONDITIONS**

Store in a refrigerator  
Do not freeze  
Keep the vial in the outer carton in order to protect from light

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE****11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Roche Registration GmbH  
Emil-Barell-Strasse 1  
79639 Grenzach-Wyhlen  
Germany

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/19/1388/002

**13. BATCH NUMBER**

Batch

**14. GENERAL CLASSIFICATION FOR SUPPLY****15. INSTRUCTIONS ON USE****16. INFORMATION IN BRAILLE**

Justification for not including Braille accepted

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

**MINIMUM PARTICULARS TO APPEAR ON SMALL IMMEDIATE PACKAGING UNITS**

**VIAL**

**1. NAME OF THE MEDICINAL PRODUCT AND ROUTE OF ADMINISTRATION**

Polivy 30 mg powder for concentrate  
polatuzumab vedotin  
Intravenous use

**2. METHOD OF ADMINISTRATION**

IV after reconstitution and dilution

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT**

30 mg

**6. OTHER**

Cytotoxic

**B. PACKAGE LEAFLET**

## Package leaflet: Information for the patient

### Polivy 30 mg powder for concentrate for solution for infusion Polivy 140 mg powder for concentrate for solution for infusion polatuzumab vedotin

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

**Read all of this leaflet carefully before you are given this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or nurse.
- If you get any side effects, talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet

1. What Polivy is and what it is used for
2. What you need to know before you are given Polivy
3. How Polivy is given
4. Possible side effects
5. How to store Polivy
6. Contents of the pack and other information

#### 1. What Polivy is and what it is used for

##### What Polivy is

Polivy is a cancer medicine that contains the active substance “polatuzumab vedotin”.

It is always used together with other cancer medicines – see below “What other medicines is Polivy given with”.

##### What Polivy is used for

Polivy is given to treat “diffuse large B-cell lymphoma” that has never been treated before.

Polivy is also given to treat “diffuse large B-cell lymphoma” that has come back or has not got better:

- after at least one previous therapy, and
- when you cannot receive a stem cell transplant.

“Diffuse large B-cell lymphoma” is a cancer that comes from “B lymphocytes” also called B-cells. These are a type of blood cells.

##### How Polivy works

Polivy contains something called a ‘monoclonal antibody’ and a substance that can kill cancer called ‘MMAE’.

- The “monoclonal antibody” part of the medicine attaches to a target on B cells.
- Once attached to B cells, the medicine releases “MMAE” into the B cells and kills them.

## **What other medicines Polivy is given with**

Polivy is given in combination with other cancer medicines:

- rituximab, cyclophosphamide, doxorubicin and prednisone for “diffuse large B-cell lymphoma” that has never been treated before.
- rituximab and bendamustine for “diffuse large B-cell lymphoma” that has come back or has not got better, after at least one previous therapy – and when you cannot receive a stem cell transplant.

## **2. What you need to know before you are given Polivy**

### **You must not be given Polivy**

- if you are allergic to polatuzumab vedotin or any of the other ingredients of this medicine (listed in section 6).
- if you currently have an active severe infection.

If the above applies to you, you must not be given Polivy. If you are not sure, talk to your doctor or nurse before you are given Polivy.

### **Warnings and precautions**

Talk to your doctor or nurse before you are given Polivy if any of the following apply to you (or you are not sure):

- you have ever had brain or nerve problems such as:
  - memory problems
  - difficulty moving or sensations in your body such as feeling pins and needles, burning, pain and discomfort even from slight touch
  - eyesight problems
- you have ever had liver problems
- you think you have an infection or have had long-lasting or repeated infections such as herpes (see “Infections” in section 4)
- you are due to have a vaccine or you know you may need to have one in the near future

If any of the above apply to you (or you are not sure) talk to your doctor or nurse before you are given Polivy.

### **Pay attention to the following side effects**

Polivy can cause some serious side effects that you need to tell your doctor or nurse about straight away. These include:

#### **Myelosuppression**

Myelosuppression is a condition in which the production of blood cells is decreased, resulting in fewer red blood cells, white blood cells, and platelets. Your doctor will do blood tests to check your blood cell count.

Tell your doctor or nurse straight away if you:

- develop chills or shivering
- have a fever
- have headaches
- feel tired
- feel dizzy
- look pale

- have unusual bleeding, bruising under the skin, bleeding longer than usual after your blood has been drawn, or bleeding from your gums.

### **Peripheral neuropathy**

Tell your doctor or nurse straight away if you have any problems with a change in the sensitivity of your skin, especially in your hands or feet, such as:

- numbness
- tingling
- a burning sensation
- pain
- discomfort or weakness
- difficulty walking.

If you had any of these symptoms before treatment with Polivy, tell your doctor straight away if you notice any changes in them.

If you have symptoms of peripheral neuropathy, your doctor may lower your dose.

### **Infections**

Signs and symptoms of infections vary between individuals, tell your doctor or nurse straight away if you develop symptoms of an infection such as:

- fever
- cough
- chest pain
- tiredness
- painful rash
- sore throat
- burning pain when passing urine
- feeling weak or generally unwell.

### **Progressive multifocal leukoencephalopathy (PML)**

PML is a very rare and life threatening infection in the brain, that has occurred in one patient treated with Polivy together with bendamustine and another medicine called obinutuzumab.

Tell your doctor or nurse straight away if you have:

- memory loss
- trouble speaking
- difficulty walking
- problems with your eyesight.

If you had any of these symptoms before treatment with Polivy, tell your doctor straight away if you notice any changes in them. You may need medical treatment.

### **Tumour lysis syndrome**

Some people may develop unusual levels of some substances (such as potassium and uric acid) in the blood caused by the fast breakdown of cancer cells during treatment. This is called “tumour lysis syndrome”. Your doctor, pharmacist or nurse will do blood tests to check for the condition.

### **Infusion-related reactions**

Infusion-related reactions, allergic or anaphylactic (more severe allergic) reactions can happen. Your doctor or nurse will check for side effects during your infusion and for 30 to 90 minutes afterwards. If you get any serious reaction, your doctor may stop treatment with Polivy.

### **Liver damage**

This medicine can cause inflammation or damage to cells in the liver that affect the normal function of the liver. Injured liver cells may leak high amounts of certain substances (liver enzymes and bilirubin) into the bloodstream, which can be detected by blood tests.

In most cases you will not have any symptoms but tell your doctor or nurse straight away if you get:

- yellowing of your skin and of the whites of your eyes (jaundice).

Your doctor will check your blood to test your liver function before and regularly during treatment.

### **Children and adolescents**

This medicine should not be used in children or young people under the age of 18. This is because there is no information about its use in this age group.

### **Other medicines and Polivy**

#### **Other medicines and vaccines**

Tell your doctor or nurse if you are taking, have recently taken or might start taking any other medicines. This includes medicines obtained without a prescription and herbal medicines.

Also tell your doctor or nurse if you are due to have a vaccine or you know you may need to have one in the near future.

### **Contraception (women and men)**

If you are a woman of childbearing age, you must use effective contraception during treatment – and for 9 months after the last dose of Polivy.

Men must use contraception during treatment – and for 6 months after the last dose of Polivy.

### **Pregnancy**

It is important to tell your doctor before and during treatment if you are pregnant, think you may be pregnant, or are planning to get pregnant. This is because Polivy can affect your baby's health.

Do not use this medicine if you are pregnant unless you and your doctor decide that the benefit to you outweighs possible risk to the unborn baby.

### **Breast-feeding**

Do not breast-feed while receiving Polivy – and for at least 3 months after the last dose, because small amount of Polivy may pass into your breast milk.

### **Fertility**

Men are advised to have sperm samples preserved and stored before treatment with this medicine.

## **Driving and using machines**

Polivy has a minor influence on your ability to drive, cycle or use any tools or machines.

- If you get infusion-related reactions or nerve damage, or if you feel tired, weak or dizzy (see section 4) do not drive, cycle or use tools or machines until the reaction stops.

See section 4 for more information about side effects.

## **Polivy contains sodium**

This medicine contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

## **3. How Polivy is given**

Polivy is given under the supervision of a doctor experienced in giving such treatments.

It is given into a vein, as a drip over 90 minutes.

### **How much Polivy is given**

The dose of this medicine depends on your body weight.

- The usual starting dose is 1.8 mg for each kilogram of your body weight.
- If you have peripheral neuropathy, your doctor may lower your dose.

### **How often is Polivy given?**

- Each cycle lasts 21 days.
- You will be given 6 treatment cycles of Polivy in combination with other medicines.

### **What other medicines is Polivy given with?**

- rituximab, cyclophosphamide, doxorubicin and prednisone for "diffuse large B-cell lymphoma" that has never been treated before or,
- rituximab and bendamustine for "diffuse large B-cell lymphoma" that has come back or has not got better, after at least one previous therapy – and when you cannot receive a stem cell transplant.

### **If you miss a dose of Polivy**

- If you miss an appointment, make another one straight away.
- For the treatment to be fully effective, it is very important not to miss a dose.

### **If you stop receiving Polivy**

Do not stop treatment with Polivy unless you have discussed this with your doctor. This is because stopping treatment may make your condition worse.

If you have any further questions on the use of this medicine, ask your doctor or nurse.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them. The following side effects have been reported with this medicine:



## Serious side effects

Tell your doctor or nurse straight away if you notice any of the following serious side effects – you may need urgent medical treatment. These may be new symptoms or a change in your current symptoms.

- infusion-related reactions – your doctor will check for these for 30-90 minutes afterwards
- fevers and chills
- rash/hives
- severe infections
- pneumonia (lung infection)
- herpes infection
- viral infections
- upper respiratory tract infection
- skin infection
- urinary tract infection
- unusual bleeding or bruising under the skin
- memory loss, trouble speaking, difficulty walking or problems with your eyesight
- yellowing of skin or whites of your eyes
- breathlessness and difficulty in breathing

## Other side effects

Tell your doctor or nurse if you notice any of the following side effects:

**Very common:** may affect more than 1 in 10 people

- pneumonia (lung infection)
- runny nose, sneezing, sore throat and cough (upper respiratory tract infection)
- numbness, tingling, a burning sensation, pain, discomfort or weakness and/or difficulty walking (peripheral neuropathy)
- fever
- cough
- vomiting
- diarrhoea or constipation
- soreness or inflammation of the mouth and/or gut (mucositis)
- feeling sick (nausea)
- abdominal (belly) pain
- feeling tired
- not feeling hungry
- loss of weight
- infusion-related reactions
- common cold
- hair loss
- changes in blood tests:
  - low levels of all types of white blood cell (combined)
  - low levels of neutrophils (a type of white blood cell) with or without fever
  - low level of platelets (a type of blood cell that helps your blood to clot)
  - low levels of red blood cells (anaemia)
  - low level of potassium in the blood (hypokalaemia)

**Common:** may affect up to 1 in 10 people

- severe infection (sepsis)
- urinary tract infection
- viral infections
- herpes infection

- skin infections
- inflammation of the lungs
- breathlessness and difficulty in breathing
- dizziness
- fluid retention causing swelling in the lower legs or hands (oedema peripheral)
- high level of transaminases in the blood
- joint pain
- itchiness
- chills
- rash
- dry skin
- muscle pain
- changes shown in blood tests:
  - decreased number of all blood cells (pancytopenia)
  - low levels of lymphocytes (a type of white blood cells)
  - low level of phosphate in the blood (hypophosphataemia)
  - low level of calcium in the blood (hypocalcaemia)
  - low level of albumin in the blood (hypoalbuminemia)
  - high level of lipase enzyme in the blood

**Uncommon: may affect up to 1 in 100 people**

- blurred vision

Tell your doctor or nurse straight away if you notice any of the side effects listed above.

**Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

**5. How to store Polivy**

Polivy will be stored by the healthcare professionals at the hospital or clinic. The storage details are as follows:

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the carton and the vial after EXP. The expiry date refers to the last day of that month.
- Store in a refrigerator (2 °C–8 °C).
- Do not freeze.
- Keep the container in the outer carton in order to protect from light.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

### What Polivy contains

- The active substance is polatuzumab vedotin.
- Polivy 30 mg: Each vial contains 30 milligrams (mg) polatuzumab vedotin.
- Polivy 140 mg: Each vial contains 140 milligrams (mg) polatuzumab vedotin.
- After reconstitution each millilitre (mL) contains 20 mg polatuzumab vedotin.
- The other ingredients are: succinic acid, sodium hydroxide, sucrose, polysorbate 20. See section “Polivy contains sodium”.

### What Polivy looks like and contents of the pack

Polivy powder for concentrate for solution for infusion is a white to slightly greyish-white cake provided in a glass vial.

Each pack of Polivy consists of one vial.

### Marketing Authorisation Holder

Roche Registration GmbH  
Emil-Barell-Strasse 1  
79639 Grenzach-Wyhlen  
Germany

### Manufacturer

Roche Pharma AG  
Emil-Barell-Strasse 1  
79639 Grenzach-Wyhlen  
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

#### **België/Belgique/Belgien**

N.V. Roche S.A.  
Tél/Tel: +32 (0) 2 525 82 11

#### **Lietuva**

UAB “Roche Lietuva”  
Tel: +370 5 2546799

#### **България**

Рош България ЕООД  
Тел: +359 2 818 44 44

#### **Luxembourg/Luxemburg**

(Voir/siehe Belgique/Belgien)

#### **Česká republika**

Roche s. r. o.  
Tel: +420 - 2 20382111

#### **Magyarország**

Roche (Magyarország) Kft.  
Tel: +36 - 1 279 4500

#### **Danmark**

Roche a/s  
Tlf: +45 - 36 39 99 99

#### **Malta**

Irreferi għall-Irlanda

#### **Deutschland**

Roche Pharma AG  
Tel: +49 (0) 7624 140

#### **Nederland**

Roche Nederland B.V.  
Tel: +31 (0) 348 438050

**Eesti**

Roche Eesti OÜ  
Tel: + 372 - 6 177 380

**Ελλάδα**

Roche (Hellas) A.E.  
Τηλ: +30 210 61 66 100

**España**

Roche Farma S.A.  
Tel: +34 - 91 324 81 00

**France**

Roche  
Tél: +33 (0) 1 47 61 40 00

**Hrvatska**

Roche d.o.o.  
Tel: +385 1 4722 333

**Ireland**

Roche Products (Ireland) Ltd.  
Tel: +353 (0) 1 469 0700

**Ísland**

Roche a/s  
c/o Icepharma hf  
Sími: +354 540 8000

**Italia**

Roche S.p.A.  
Tel: +39 - 039 2471

**Κύπρος**

Γ.Α.Σταμάτης & Σια Λτδ.  
Τηλ: +357 - 22 76 62 76

**Latvija**

Roche Latvija SIA  
Tel: +371 - 6 7039831

**Norge**

Roche Norge AS  
Tlf: +47 - 22 78 90 00

**Österreich**

Roche Austria GmbH  
Tel: +43 (0) 1 27739

**Polska**

Roche Polska Sp.z o.o.  
Tel: +48 - 22 345 18 88

**Portugal**

Roche Farmacêutica Química, Lda  
Tel: +351 - 21 425 70 00

**România**

Roche România S.R.L.  
Tel: +40 21 206 47 01

**Slovenija**

Roche farmacevtska družba d.o.o.  
Tel: +386 - 1 360 26 00

**Slovenská republika**

Roche Slovensko, s.r.o.  
Tel: +421 - 2 52638201

**Suomi/Finland**

Roche Oy  
Puh/Tel: +358 (0) 10 554 500

**Sverige**

Roche AB  
Tel: +46 (0) 8 726 1200

**United Kingdom (Northern Ireland)**

Roche Products (Ireland) Ltd.  
Tel: +44 (0) 1707 366000

**This leaflet was last revised in {MM/YYYY}**

**Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu/>

-----  
The following information is intended for healthcare professionals only:

Procedures for proper handling and disposal of anticancer medicinal products should be considered.

#### Instructions for reconstitution

- Polivy 30 mg: Using a sterile syringe, slowly inject 1.8 mL of sterile water for injection into the 30 mg Polivy vial to yield a single-dose solution containing 20 mg/mL polatuzumab vedotin. Direct the stream toward the wall of the vial and not directly on the lyophilized cake.
- Polivy 140 mg: Using a sterile syringe, slowly inject 7.2 mL of sterile water for injection into the 140 mg Polivy vial to yield a single-dose solution containing 20 mg/mL polatuzumab vedotin. Direct the stream toward the wall of the vial and not directly on the lyophilized cake.
- Swirl the vial gently until completely dissolved. Do not shake.
- Inspect the reconstituted solution for discoloration and particulate matter. The reconstituted solution should appear colourless to slightly brown, clear to slightly opalescent, and free of visible particulates. Do not use if the reconstituted solution is discoloured, is cloudy, or contains visible particulates.

#### Instructions for dilution

1. Polivy must be diluted to a final concentration of 0.72-2.7 mg/mL in an intravenous infusion bag, with a minimum volume of 50 mL, containing 9 mg/mL sodium chloride solution for injection, or 4.5 mg/mL sodium chloride solution for injection, or 5% glucose.
2. Determine the volume of 20 mg/mL reconstituted solution needed based on the required dose (see below):  
Total Polivy dose (mL) to be further diluted = 
$$\frac{\text{Polivy dose (mg/kg)} \times \text{patient's weight (kg)}}{\text{Reconstituted vial concentration (20 mg/mL)}}$$
3. Withdraw the required volume of reconstituted solution from the Polivy vial using a sterile syringe and dilute into the intravenous infusion bag. Discard any unused portion left in the vial.
4. Gently mix the intravenous bag by slowly inverting the bag. Do not shake.
5. Inspect the intravenous bag for particulates and discard if present.

#### Reconstituted solution

From a microbiological point of view, the reconstituted solution should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours refrigerated (2 °C–8 °C), unless reconstitution has taken place in controlled and validated aseptic conditions. Chemical and physical in-use stability of the reconstituted solution has been demonstrated for up to 72 hours refrigerated (2 °C–8 °C) and up to 24 hours at room temperature (9 °C–25 °C).

#### Diluted solution

From a microbiological point of view, the prepared solution for infusion should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours refrigerated (2 °C–8 °C), unless dilution has taken place in controlled and validated aseptic conditions. Chemical and physical stability of the prepared solution for infusion has been demonstrated for the durations listed in Table 1. Discard diluted Polivy solution if storage time exceeds the limits specified in Table 1.

**Table 1 Durations for which chemical and physical stability of the prepared solution for infusion have been demonstrated**

| <b>Diluent used to prepare solution for infusion</b> | <b>Solution for infusion storage conditions<sup>1</sup></b>                                 |
|--|---|
| Sodium chloride 9 mg/mL (0.9%)                       | Up to 72 hours refrigerated (2 °C–8 °C) or up to 4 hours at room temperature (9 °C–25 °C)   |
| Sodium chloride 4.5 mg/mL (0.45%)                    | Up to 72 hours refrigerated (2 °C–8 °C) or up to 8 hours at room temperature (9 °C–25 °C)   |
| 5% glucose   | Up to 72 hours refrigerated (2 °C – 8 °C) or up to 8 hours at room temperature (9 °C–25 °C) |

<sup>1</sup> To ensure product stability, do not exceed specified storage durations.